
Force transmission is a master regulator of mechanical cell competition

In the format provided by the
authors and unedited

The PDF file includes:

Supplementary Video captions 1 to 9
Supplementary Figures 1 to 10

Other Supplementary Materials for this manuscript include the following:

Supplementary Videos 1 to 9
Source data files

Video captions:

Video S1: Timelapse video of competition within patient-derived metaplastic cancer cells. Phase contrast images of E-cad⁺ cells surrounded by E-cad⁻ cells corresponding to **Figure 1B**. Frame rate 10 min. Scale bar 200 μ m.

Video S2: Timelapse video of competition within patient-derived metaplastic cancer cells derived from a second patient. Phase contrast images of E-cad⁺ cells surrounded by E-cad⁻ cells corresponding to **Fig. S1 C**. Frame rate 10 min. Scale bar 200 μ m.

Video S3: Timelapse video of competition between MDCK WT and MDCK E-cad KO cells. Phase contrast images of E-cad KO cells expressing LifeAct-EGFP in green. Frame rate 15 min. Scale bar 100 μ m.

Video S4: Timelapse video of competition between MDCK dKO and MDCK E-cad KO cells. Phase contrast images of E-cad KO cells expressing LifeAct-EGFP in green. Frame rate 15 min. Scale bar 100 μ m.


Video S5: Laser ablation of WT cluster within the mixed culture. WT cells express CAAX-GFP and E-cad KO cells express LifeAct-EGFP. Video corresponds to Fig. S5 D. Magenta line shows ablation area. Frame rate 5 sec. Scale bar 50 μ m.

Video S6: Laser ablation of E-cad KO cluster within the mixed culture. WT cells express CAAX-GFP and E-cad KO cells express LifeAct-EGFP. Video corresponds to Fig. S5 D. Magenta line shows ablation area. Frame rate 5 sec. Scale bar 50 μ m.


Video S7: Timelapse video of cell eliminations at the interface between MDCK WT and MDCK E-cad KO cells. Phase contrast images of E-cad KO expressing LifeAct-EGFP in green. Magenta line highlights the interface. Note that E-cad KO cells are not only eliminated when directly in contact with WT cells. Frame rate 15 min. Scale bar 50 μ m.

Video S8: Example video showing simulation of competition between model WT and model E-cad KO cells. mWT cells are shown in blue, mE-cad KO cells in red. The cell-cell adhesion strength of mWT cells is 8x higher than the mE-cad KO, the cell-substrate adhesion strength is constant.


Video S9: Timelapse video of actin dynamics in competition between MDCK WT and E-cad KO cells. Confocal images showing maximum projection of actin expressed in both cell types. 60% MDCK WT cells express LifeAct-mRuby, displayed in white, all E-cad KO cells express LifeAct-EGFP, displayed in green. Note the protrusion activity of E-cad KO cells at the interface. Frame rate 7.5 min. Scale bar 20 μ m.




Supplementary Figure 1: E-cad⁺ triple negative tumor cells are viable and sample from second patient (A) Upper: Mix of both sub-populations. The E-cad⁺ population is spreading, leading to eliminations of E-cad⁻ cells in its vicinity. Lower: E-cad⁻ cells grow to high densities independent of interactions with E-cad⁺ cells. Over time, elimination at random positions can be observed, but fewer than when interacting with the E-cad⁺ sub-population. (B) Confocal image of metaplastic breast cancer cells derived from a second patient stained for E-cadherin (magenta). E-cad⁺ and E-cad⁻ cells have sorted. (C) Phase contrast images of cells corresponding to (B). Magenta line shows initial boundaries of the sub-populations red lines show boundaries at the given timepoint. Representative images from N=3 independent experiments. Scale bars 200 μ m.




Supplementary Figure 2: E-cad KO cells lose independent of ratio or level of sorting (A) Confocal image of nuclei (white) and E-cad KO cells (green). Middle: Corresponding nuclei segmentation using StarDist. The cellular identity was determined based on the E-cad KO fluorescence. Right: Average cell densities in mixed cultures. Each datapoint shows the cell density for one FOV. $n=16$ FOV from $N=2$ independent experiments. P-value from unpaired, two-sided t-test. **(B)** Left: Snapshots from Airyscan confocal movies of WT cells expressing LifeAct-mCherry and E-cad KO cells expressing LifeAct-EGFP. The E-cad KO cells are eliminated by LifeAct expressing WT cells. Right: Area quantifications corresponding to images on the left from $n=4$ movies from $N=1$ experiment. **(C)** Brightfield and fluorescence images of WT and E-cad KO cells (green) at the beginning of the experiment and 12 and 24 hours later. Magenta lines show initial cluster sizes to visualize cluster progression over time. Bottom row: Area development of E-cad KO cells. (i) E-cad KO in minority. (ii) E-cad KO in majority. (iii) Collision of E-cad KO and WT tissues. Movies are analysed shortly after contact formation. Area quantifications from (i) $n=8$ movies from $N=3$ independent experiments, (ii) $n=7$ movies from $N=3$ independent experiments, (iii) $n=4$ movies from $N=2$ independent experiments. Data are presented as mean values \pm SD. Scale bars 200 μm (A), 100 μm (C), 10 μm (B).




Supplementary Figure 3: Mechanics of intratumoral cell competition (A) Brightfield image of E-cad (-) and E-cad (+) tumor clusters, as shown in **Figure 1**. (B) Cell stiffnesses measured with nanoindentation. Each datapoint shows one measurement. N=2 independent experiments and n=57 indentations. (C) Map of traction forces corresponding to (A). Green lines show cluster outlines. Right: Average traction forces. Each datapoint shows the average value from one field-of-view at one timepoint. N=2 independent experiments and n=408 timepoints. P-values from two-sided t-test. Data are presented as mean values +/- SD. Scale bars 100 μ m.




Supplementary Figure 4: Mixed culture with low cell-substrate adhesion (A) Confocal images of focal adhesions (Paxillin, white) in mixed culture grown on (i) glass or on (ii) soft substrate (370 Pa, PAA). **(B)** Left: map of traction forces on soft substrate. Right: Average traction forces on different substrate stiffnesses. Green lines show outlines of E-cad KO cluster. Each datapoint represents one average map. $n=568$ positions from $N=4$ independent experiments (15 kPa) and $n=170$, $N=2$ (370 Pa). P-values from Kruskal-Wallis test corrected for multiple comparisons (Dunn's test). Data are presented as mean values \pm SD. Scale bars 20 μ m.




Supplementary Figure 5: Actomyosin accumulation at interface between E-cad KO and WT cells and interface elimination during intratumoral competition (A) Representative airyscan confocal image of increased phospho-myosin accumulating in both cell types at the interface. Intensity line plots were drawn at the indicated position for actin (phalloidin), phospho-myosin and LifeAct-EGFP (E-cad KO cells). Zoom-in on intensity profile reveals two separate intensity peaks in WT and E-cad KO and the colocalization of LifeAct-EGFP and phospho-myosin. (B) Actin dynamics following cell-cell collision. WT cells express LifeAct-mCherry (white) and E-cad KO cells express LifeAct-EGFP (green). Intensity values were measured at the indicated line. An increase in actin intensity within the WT cells was observed following collision (red arrow) at new interface with E-cad KO cells. (C) Representative phase-contrast image of patient-derived tumor xenograft cultured in 2D. E-cad (-) cells are eliminated at the interface. (D) Confocal images showing an increase of phospho-myosin at the tissue interface. Scale bars 200 μm (C), 20 μm (A), 25 μm (D), 10 μm (B).




Supplementary Figure 6: Preferred interface elimination is independent of interface curvature Brightfield and fluorescence (E-cad KO) images of a competing MDCK coculture. Within the same field of view, E-cad KO cells are eliminated at positively (i), minimally (ii) or negatively (iii) curved interfaces (dashed lines). Extrusions indicated by red circles. Scale bars: 100 μm (overview), 10 μm (zoom-ins).




Supplementary Figure 7: Schematic of shape changes The analytical model uses these three shapes: a cylinder, a cone and a sphere, to calculate the change in area used to find the work. All three cell shapes have the same volume.



Supplementary Figure 8: Key features of the multi-phase field model An example cross-section from a simulation. The arrows show, schematically, how cell-cell and cell-substrate adhesions are explicitly accounted for in the model and mathematically outlined in the Methods.



Supplementary Figure 9: Stress fluctuations away from the interface highlight the role of stress transmission Susceptibility of two-dimensional isotropic stress (σ_{2D}^{iso}), and the out-of plane stress component (σ_{zz}) for mE-cad KO cells as a function of distance away from the interface for various cell-substrate adhesion ($\tilde{\omega}_{cs}$) and cell-cell adhesion ($\tilde{\omega}_{cc}$) strengths. The distance is normalized by the initialized cell radius. The susceptibility is normalized by the maximum value.



Supplementary Figure 10: Phase contrast and fluorescence images of 15 kPa PDMS coated with Cy3-labelled fibronectin. Left: Before cell seeding, right: after 3 days of cell culture. Scale bars 200 μm .